

## Skin (cutaneous) B-cell lymphoma

Skin (cutaneous) lymphomas are non-Hodgkin lymphomas that start in the skin. This page is about a rare type of skin lymphoma called B-cell skin lymphoma (also known as cutaneous B-cell lymphoma or CBCL).

### On this page

[What is B-cell skin lymphoma?](#)

[Who gets it?](#)

[Symptoms](#)

[Diagnosis and staging](#)

[Types of B-cell skin lymphoma](#)

[Low-grade B-cell skin lymphoma](#)

[High-grade B-cell skin lymphoma](#)

[Outlook](#)

[Follow-up](#)

[Research and clinical trials](#)

## What is B-cell skin lymphoma?

**Lymphoma** is a cancer that starts in white blood cells called lymphocytes, which are part of your **immune system**. There are two types of lymphocyte: **T lymphocytes (T cells)** and **B lymphocytes (B cells)**. Lymphomas can develop from either T cells or B cells.

Skin lymphomas are lymphomas that develop in the skin and have not affected any other areas of the body at the time they are diagnosed. B-cell skin lymphomas are skin lymphomas that develop from B cells.

A lymphoma that starts somewhere else in the body and then spreads to the skin is **not** a skin lymphoma. If you have a lymphoma that has spread to the skin, our information on the particular **type of lymphoma** you have will be more relevant for you.

We have separate information on **T-cell skin lymphomas** (skin lymphomas that develop from T cells).

---

## Who gets B-cell skin lymphoma?

B-cell skin lymphoma (also called cutaneous B-cell lymphoma or CBCL) is rare. Only around 100 people are diagnosed with B-cell skin lymphoma in the UK each year.

Scientists don't know exactly what **causes** skin lymphoma but there are some factors that might increase your risk of developing it. These vary according to the type of skin lymphoma.

---

## Symptoms of B-cell skin lymphoma

The main symptom of B-cell skin lymphoma is a lump or lumps in the skin. The lumps often appear as small, raised, solid areas of skin that can look like small pimples. These are called papules. They may develop into thickened but still quite flat areas called plaques, or larger lumps, often deep red or purplish, called nodules or tumours. Papules or nodules appear most commonly on the head, neck, back or legs.

**It is important to remember that B-cell skin lymphomas are rare. There are many skin conditions that may look similar but are not lymphomas.**



**Figure: A tumour of B-cell skin lymphoma on the head**

You might have one lump or plaque, or you might have several, which may be grouped together or more widely spread out. Sometimes the lumps break down (ulcerate) and may become infected.

For more information about these symptoms, including photographs, see our [overview of skin lymphomas](#).

You may also have [swollen lymph nodes](#) in your neck, armpits or groin. They may be inflamed as a reaction to the nearby skin irritation or because they contain abnormal lymphoma cells.

A few people experience more general [symptoms](#) of lymphoma such as unexplained weight loss, fevers or night sweats. However, this is quite uncommon with B-cell skin lymphoma.

Many skin lymphomas develop very slowly, over months or years. They can look like more common skin conditions, such as eczema or psoriasis, or reactions to insect bites or vaccinations. Because of this, it may take a long time for your GP to rule out other conditions and refer you to a specialist. This might be a specialist in skin diseases (dermatologist) or a specialist in diseases of the blood and **lymphatic system** (haematologist).

---

## Diagnosis and staging of B-cell skin lymphoma

B-cell skin lymphomas are diagnosed using a **skin biopsy**. A skin biopsy can find out the exact type of skin lymphoma you have and whether it is slow-growing (**low-grade** or indolent) or fast-growing (**high-grade** or aggressive). However, diagnosing skin lymphoma is not always straightforward, even for a specialist. You might need several skin biopsies before your doctor is able to confirm your diagnosis.

You may also have other tests to find out if any other parts of your body are affected by lymphoma – this is called '**staging**'. Other tests may include:

- a physical examination
- **blood tests** to look at your general health and to test for bacterial or viral infections that might be linked to your lymphoma
- a **CT scan** or **PET/CT scan**
- a **bone marrow biopsy**
- a **lymph node biopsy** (if you have swollen lymph nodes).

**Waiting for the results** of your tests can be difficult but it is important that your specialist knows exactly what type of lymphoma you have so they can give you the most appropriate **treatment**.

## Staging of B-cell skin lymphoma

The stage of the lymphoma describes how far it has grown. The stage, along with the particular type of lymphoma you have, helps your medical team decide on the most appropriate treatment for you. For B-cell skin lymphomas, the stage of your lymphoma when it is diagnosed does **not** affect your outlook (prognosis).

If your lymphoma is only affecting your skin, it might be staged as a lymphoma that has started outside the **lymphatic system** (an 'extranodal' lymphoma) but has not spread anywhere else. This is known as **Stage 1E**. Doctors might add an 'A' after it (Stage 1AE). This means you do not have any symptoms known as 'B symptoms' (unexplained weight loss, fevers or night sweats).

Otherwise, B-cell skin lymphomas are staged using a system called 'TNM'. TNM stands for:

- **Tumour:** how many papules, nodules or tumours you have, how big they are and where they are. This is shown by a 'T' and a number between 1 and 3, sometimes followed by a letter between a and c.
- **Nodes:** how many lymph nodes are involved (if any) and where they are. This is shown by an 'N' and a number between 0 and 3.
- **Metastasis:** whether the lymphoma has spread (metastasised) to any part of the body apart from your skin or lymph nodes. This is shown by an 'M' followed by 0 or 1.

Your stage includes all three measures. For example:

- a person with one skin nodule smaller than 5cm, no swollen lymph nodes and no lymphoma anywhere else in their body would have a stage of T1aN0M0
- a person with several skin nodules confined to a 30cm area of skin, swollen lymph nodes in two areas of their body but no lymphoma anywhere else would have a stage of T2bN2M0.

Staging helps monitor changes in your lymphoma over time. This can help your doctors decide on the best treatment for you.

---

## Types of B-cell skin lymphoma

B-cell skin lymphomas have technical names based on what the lymphoma cells look like under a microscope. The different types behave differently and may be treated differently depending on whether they are slow-growing (**low-grade**) or fast-growing (**high-grade**). There are three main types:

- primary cutaneous follicle centre lymphoma (low-grade)
  - primary cutaneous marginal zone lymphoma (low-grade)
  - primary cutaneous diffuse large B-cell lymphoma, leg-type (high-grade).
-

## Low-grade B-cell skin lymphomas

Primary cutaneous follicle centre lymphoma is the most common type of B-cell skin lymphoma. It accounts for over half of all B-cell skin lymphomas. It is a slow-growing (low-grade) lymphoma that develops over months or years. It typically develops in people in their 50s and is slightly more common in men than women. It usually appears as pink, reddish or purple papules, plaques or nodules, most commonly on the head, neck, chest or back. It often develops as a single lesion but you might have several lesions, usually grouped together.

Primary cutaneous marginal zone lymphoma is another low-grade lymphoma. It typically develops in young adults (35 to 60 years old) and is slightly more common in men than women. Some cases are linked to an infection with a bacterium called *Borrelia burgdorferi*. This is carried by ticks and causes Lyme disease. However, most people with Lyme disease do not develop skin lymphoma.

Primary cutaneous marginal zone lymphoma is also known as '**extranodal marginal zone lymphoma of mucosa-associated lymphoid tissue**' (MALT lymphoma). It usually appears as pink, red or purplish papules, plaques or nodules that develop on your arms, chest or back, or sometimes your legs. They can develop in one place or in several places and are very slow-growing.

### Treatment of low-grade B-cell skin lymphomas

If you have primary cutaneous marginal zone lymphoma and your blood tests show you have an infection with *Borrelia burgdorferi*, you have treatment with antibiotics first. This might be the only treatment you need.

Otherwise, all types of low-grade B-cell skin lymphoma are treated in the same way.

- If only one, small area of skin is affected by lymphoma, you might have surgery to remove it. This may be followed by radiotherapy to prevent it coming back (relapsing).
- If just a few areas of your skin are affected, you usually have radiotherapy on its own. Surgery is not usually used for more than one site.

If your lymphoma is more widespread, localised radiotherapy or surgery are generally not suitable. Treatment depends on how much of your skin is affected and whether the lymphoma has spread anywhere else.

- If you have lymphoma in several areas of your skin but you are not experiencing any troublesome symptoms, you might not need treatment straightaway. Instead, your doctor might monitor your progress. This is called **active monitoring or watch and wait**. It does not mean there is no treatment for you, but that you would not benefit from starting treatment straightaway. This saves you experiencing side effects of unnecessary treatment. You have treatment when you need it.
- If you have lymphoma in several areas of your skin and you are experiencing symptoms, you might have a **chemotherapy** drug called chlorambucil or an **antibody treatment** called **rituximab**. This can be given through a drip into a vein (intravenously) or as an injection directly into the affected skin.
- If your lymphoma has spread to your lymph nodes or other parts of your body, you might have rituximab plus a **chemotherapy regimen** (combination of treatment) such as bendamustine or CVP (cyclophosphamide, vincristine and prednisolone).

If your lymphoma comes back (**relapses**) after treatment, it can often be successfully treated with the same treatment you had the first time.

There is more information on our page on **treatment of skin lymphoma**.

---

## High-grade B-cell skin lymphomas

Primary cutaneous diffuse large B-cell lymphoma, leg-type, is a very rare, fast-growing (high-grade) type of lymphoma. It typically develops in people over 70 and is twice as common in women as men. It usually develops as one or more large, bluish-red **nodules** that grow rapidly over a few weeks or months. They are usually on the lower legs but around 1 in 10 develop in other places. They can become quite large and deep and they may break down (ulcerate).

### Treatment of high-grade B-cell skin lymphomas

Primary cutaneous diffuse large B-cell lymphoma, leg type, is usually treated with **chemotherapy**, often followed by **radiotherapy** to the affected area. The **chemotherapy regimen** (combination of treatment) most commonly used is called CHOP (cyclophosphamide, hydroxydaunorubicin, vincristine [also known as **Oncovin**®] and prednisolone). It is usually given along with an **antibody treatment** called **rituximab** (R-CHOP).

If you are not well enough to have chemotherapy, you might have radiotherapy on its own to help control your symptoms. Some people have rituximab on its own.

If the lymphoma comes back (**relapses**), your doctor will probably suggest more chemotherapy.

There is more information on our page on [treatment of skin lymphoma](#).

---

## Outlook for B-cell skin lymphoma

Your outlook (prognosis) depends on lots of factors, including your age and general health and the exact type of skin lymphoma you have. Your specialist is best placed to advise you on your outlook based on your individual circumstances.

In general, slow-growing (**low-grade**) B-cell skin lymphomas have an excellent outlook. They are often diagnosed early, grow slowly and respond well to treatment. Any skin problems they cause come and go and only need treatment some of the time.

Some slow-growing lymphomas do not become obvious until they are more advanced and the disease has spread to the lymph nodes, blood or internal organs. With treatment, many people with advanced low-grade skin lymphoma have long periods when they are free of symptoms.

Primary cutaneous diffuse large B-cell lymphoma, leg type, is a fast-growing (**high-grade**) lymphoma and is more aggressive in its behaviour. It needs more intensive treatment more urgently. Most people respond to treatment but it is more likely to come back (relapse) than low-grade B-cell skin lymphoma.

Occasionally, slow-growing (low-grade) B-cell skin lymphomas can change (**transform**) into a faster-growing type, although this is very rare. Your medical team should check for this. If your lymphoma transforms, you need more intensive treatment.

---



## Follow-up of B-cell skin lymphoma

How often you have **check-ups** after your treatment ends depends mainly on what type of skin lymphoma you have and how it is responding to treatment. You might only see your specialist every 6 to 12 months if:

- you have a slow-growing (low-grade) skin lymphoma
- your condition is stable (unchanging) or in **remission** (no evidence of the disease) after treatment.

If your skin lymphoma is fast-growing (high-grade), or is continuing to grow, you may see your doctor every 4 to 6 weeks.

At the clinic, your doctor will ask about your symptoms and examine you. Occasionally, you might have blood tests, scans or biopsies.

**Tell your doctor if you have any symptoms that are troubling you, if you've noticed any changes in your skin, or you have any new swellings.**

---

## Research and clinical trials

Researchers are continually trying to find out which treatment or combination of treatments work best for skin lymphomas. Your doctor may ask you to consider taking part in a clinical trial. Clinical trials allow new treatments to be evaluated and compared with more established ones. Studying treatments is the only way that new and, hopefully, better treatments can become available.

Find out more about clinical trials and search for a trial that might be suitable for you at [Lymphoma TrialsLink](#).

---

We have separate information about the topics in **bold font**. Please get in touch if you'd like to request copies or if you would like further information about any aspect of lymphoma. Phone 0808 808 5555 or email [information@lymphoma-action.org.uk](mailto:information@lymphoma-action.org.uk).

## References

The full list of references for this page is available on our website. Alternatively, email [publications@lymphoma-action.org.uk](mailto:publications@lymphoma-action.org.uk) or call 01296 619409 if you would like a copy.

## Acknowledgements

- Dr Eileen Parry, Consultant Dermatologist, Tameside and Glossop Integrated Care NHS Foundation Trust. Dr Parry has been paid to take part in advisory board meetings for Actelion, Takeda and Kyowa-Kirin and has provided expert opinion on NICE submissions.
- Helen Easson, Haematology Clinical Nurse Specialist, Royal Hallamshire Hospital.
- We would like to thank the members of our Reader Panel who gave their time to review this information.
- With thanks to Kyowa Kirin for contributing towards the costs of maintaining our skin lymphomas webpages. In accordance with [how we work with healthcare and pharmaceutical companies](#), Kyowa Kirin has had no involvement in the content.

---

Content last reviewed: April 2019

Next planned review: April 2022

LYMweb0200BSkin2019v2



© Lymphoma Action

Tell us what you think and help us to improve our resources for people affected by lymphoma. If you have any feedback, please visit [lymphoma-action.org.uk/Feedback](https://lymphoma-action.org.uk/Feedback) or email [publications@lymphoma-action.org.uk](mailto:publications@lymphoma-action.org.uk).

All our information is available without charge. If you have found it useful and would like to make a donation to support our work you can do so on our website [lymphoma-action.org.uk/Donate](https://lymphoma-action.org.uk/Donate). Our information could not be produced without support from people like you. Thank you.

### Disclaimer

We make every effort to make sure that the information we provide is accurate at time of publication, but medical research is constantly changing. Our information is not a substitute for individual medical advice from a trained clinician. If you are concerned about your health, consult your doctor.

Lymphoma Action cannot accept liability for any loss or damage resulting from any inaccuracy in this information or third party information we refer to, including that on third party websites.

Neither the Information Standard scheme operator nor the scheme owner shall have any responsibility whatsoever for costs, losses or direct or indirect damages or costs arising from inaccuracy of information or omissions in the information published on the website on behalf of Lymphoma Action.